



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/10, 9/26, 15/62	A1	(11) International Publication Number: WO 97/40146 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/US96/05727 (22) International Filing Date: 24 April 1996 (24.04.96) (71) Applicant: GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US). (72) Inventor: JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). (74) Agent: BROWN, Scott, A.; Genetics Institute, Inc., 87 CambridgePark Drive, Cambridge, MA 02140 (US).		(81) Designated States: AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: YEAST INVERTASE GENE AS REPORTER SYSTEM FOR ISOLATING CYTOKINES (57) Abstract A novel method for isolating novel secreted mammalian proteins is described in which mammalian secretory leader sequences are detected using the yeast invertase gene as a reporter system.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

YEAST INVERTASE GENE AS REPORTER SYSTEM FOR ISOLATING CYTOKINES

5 The present invention relates to a novel method of cloning cDNAs which encode cytokines, to novel cDNAs isolated by the method, and to novel secreted proteins encoded by the cDNAs.

BACKGROUND OF THE INVENTION

10

Cytokines are secreted proteins which act on specific hematopoietic target cells to cause a differentiation event or on other target cells to induce a particular physiological response, such as secretion of proteins characteristic of inflammation. Cytokines, also variously known as lymphokines, hematopoietins, interleukins, colony stimulating factors, and the like, can be important therapeutic agents, especially for diseases or conditions in which a specific cell population is depleted. For example, erythropoietin, G-CSF, and GM-CSF, have all become important for treatment of anemia and leukopenia, respectively. Other cytokines such as interleukin-3, interleukin-6 and interleukin-11 show promise in treatment of conditions such as thrombocytopenia.

20

For these reasons a significant research effort has been expended in searching for novel cytokines and cloning the DNAs which encode them. In the past, novel cytokines were identified by assaying a particular cell such as a bone marrow cell, for a measurable response, such as proliferation. The search for novel cytokines has thus been limited by the assays available, and if a novel cytokine has an activity which is unmeasurable by a known assay, the cytokine remains undetectable. In a newer approach, cDNAs encoding cytokines have been detected using the polymerase chain reaction (PCR) and oligonucleotide primers having homology to shared motifs of known cytokines or their receptors. The PCR approach is also limited by the necessity for knowledge of previously cloned cytokines in the same protein family. Cytokines have also been cloned using

25

30

subtractive hybridization to construct and screen cDNA libraries, or they can potentially be cloned using PCR followed by gel electrophoresis to detect differentially expressed genes. The subtractive hybridization methods are based on the assumption that cytokine mRNAs are those that are differentially expressed, and these methods do not require any prior knowledge of the sequence of interest. However, many cytokines may be encoded by mRNAs which are not differentially expressed, and thus are undetectable using these methods.

SUMMARY OF THE INVENTION

10

In one embodiment, the present invention is directed to a method for isolating a cDNA encoding a novel secreted mammalian protein which comprises:

- a) constructing a cDNA library from mammalian cellular RNA;
- b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
- c) transforming the ligated DNA into *E. coli*;
- d) isolating plasmid DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed *E. coli* of step c);
- e) transforming the DNA of step d) into a yeast cell which does not contain the invertase gene;
- f) selecting yeast cells capable of growth on sucrose or raffinose;
- g) purifying DNA from step f) which contains a novel mammalian leader sequence ligated to yeast invertase;
- h) screening a second cDNA library to detect full-length cDNAs which contain the novel mammalian leader sequence of step g); and
- i) isolating the full-length cDNA of step h).

25

20

15

In other embodiments, the invention provides a method for screening for a cDNA encoding a novel secreted mammalian protein which comprises:

30

- a) constructing a cDNA library from mammalian cellular RNA;

- 5 b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
- c) transforming the DNA of step b) into a yeast cell which does not contain an invertase gene;
- d) selecting transformed yeast cells from step c) which are capable of growth on sucrose or raffinose;
- e) purifying DNA from the yeast cells of step d);
- 10 f) analyzing the DNA obtained from step e) to determine its sequence and to determine whether it contains a novel sequence;
- g) preparing a second cDNA library from mammalian cellular RNA and screening said second cDNA library to detect full-length cDNAs which contain the novel sequence of step f); and
- 15 h) isolating the full-length cDNA of step g) wherein the isolated cDNA encodes a putative secreted mammalian protein.

In other preferred embodiments, the method comprises the following additional steps:

- 20 1) transforming the ligated DNA into bacteria;
- 2) isolating DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed bacteria of step 1);

wherein the additional steps are performed after step b) and before step c), and wherein the DNA isolated in step 2) is used for the transformation in step c).

25 Preferably, the bacteria is *E. coli*.

In another embodiment, the invention is directed to a cDNA encoding a secreted protein isolated using the method of:

- a) constructing a cDNA library from mammalian cellular RNA;
- 30 b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;

- c) transforming the ligated DNA into E. coli;
- d) isolating plasmid DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed E. coli of step c);
- 5 e) transforming the DNA of step d) into a yeast cell which does not contain the invertase gene;
- f) selecting yeast cells capable of growth on sucrose or raffinose;
- g) purifying DNA from step f) which contains a novel mammalian leader sequence ligated to yeast invertase;
- 10 h) screening a second cDNA library to detect full-length cDNAs which contain the novel mammalian leader sequence of step g); and
- i) isolating the full-length cDNA of step h).

In other embodiments, the invention provides a method for screening for a cDNA encoding a novel secreted mammalian protein which comprises:

- 15 a) constructing a cDNA library from mammalian cellular RNA;
- b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
- c) transforming the DNA of step b) into a yeast cell which does not contain an invertase gene;
- 20 d) selecting transformed yeast cells from step c) which are capable of growth on sucrose or raffinose;
- e) purifying DNA from the yeast cells of step d);
- f) analyzing the DNA obtained from step e) to determine its sequence and to determine whether it contains a novel sequence;
- 25 g) preparing a second cDNA library from mammalian cellular RNA and screening said second cDNA library to detect full-length cDNAs which contain the novel sequence of step f); and
- 30 h) isolating the full-length cDNA of step g) wherein the isolated

cDNA encodes a putative secreted mammalian protein.

In other preferred embodiments, the method comprises the following additional steps:

- 1) transforming the ligated DNA into bacteria;
- 5 2) isolating DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed bacteria of step 1);

wherein the additional steps are performed after step b) and before step c), and wherein the DNA isolated in step 2) is used for the transformation in step c).

10 Preferably, the bacteria is *E. coli*.

In yet another embodiment the invention comprises a secreted protein isolated by the steps of:

- a) constructing a cDNA library from mammalian cellular RNA;
- b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
- 15 c) transforming the ligated DNA into *E. coli*;
- d) isolating plasmid DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed *E. coli* of step c);
- 20 e) transforming the DNA of step d) into a yeast cell which does not contain the invertase gene;
- f) selecting yeast cells capable of growth on sucrose or raffinose;
- g) purifying DNA from step f) which contains a novel mammalian leader sequence ligated to yeast invertase;
- 25 h) screening a second cDNA library to detect full-length cDNAs which contain the novel mammalian leader sequence of step g);
- i) isolating the full-length cDNA of step h);
- j) expressing the full-length cDNA of step i) in a host cell grown in a suitable culture medium; and
- 30 k) isolating the protein expressed in step j) from the culture medium.

In other embodiments, the invention provides secreted mammalian protein isolated by a method comprising:

- a) constructing a cDNA library from mammalian cellular RNA;
- b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
- c) transforming the DNA of step b) into a yeast cell which does not contain an invertase gene;
- d) selecting transformed yeast cells from step c) which are capable of growth on sucrose or raffinose;
- e) purifying DNA from the yeast cells of step d);
- f) analyzing the DNA obtained from step e) to determine its sequence and to determine whether it contains a novel sequence;
- g) preparing a second cDNA library from mammalian cellular RNA and screening said second cDNA library to detect full-length cDNAs which contain the novel sequence of step f);
- h) isolating the full-length cDNA of step g) wherein the isolated cDNA encodes a putative secreted mammalian protein;
- i) expressing the full-length cDNA of step h) in a host cell grown in a suitable culture medium; and
- j) isolating the protein expressed in step i) from the culture medium.

In other preferred embodiments, the method comprises the following additional steps:

- 1) transforming the ligated DNA into bacteria;
- 2) isolating DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed bacteria of step 1);

wherein the additional steps are performed after step b) and before step c), and wherein the DNA isolated in step 2) is used for the transformation in step c).

Preferably, the bacteria is *E. coli*.

DETAILED DESCRIPTION OF THE INVENTION

5 In order to use sucrose or raffinose as a carbon and energy source, yeast such as *Saccharomyces cerevisiae* must secrete the enzyme invertase, which cleaves sucrose to yield fructose and glucose and which cleaves raffinose to yield sucrose and melibiose. A large number of known mammalian secretory leader sequences can mediate secretion of yeast invertase. In accordance with the present invention,
10 therefore, mammalian cDNA libraries are screened first for novel secretory leader sequences which can mediate secretion of yeast invertase; and second, for the full-length cDNAs containing those novel secretory leader sequences. In this way novel secreted proteins are selected using a method which requires neither a bioassay nor knowledge of homology with other proteins.

15 Invertase genes appropriate for use in the method of the invention must encode only a nonsecreted enzyme. Preferably, the DNA encoding the invertase secretory leader sequence is removed. More preferably, the DNA encoding the invertase secretory leader sequence and the initiating methionine codon are removed. Most preferably, the DNA encoding the invertase secretory leader
20 sequence, the initiating methionine and the first two codons (for methionine and serine) of the mature invertase protein are removed. Numerous methods for selective removal of DNA segments are known.

 A nonsecreted invertase gene from any yeast species or strain may be used in the method of the invention. Preferably, the invertase gene from *Saccharomyces cerevisiae* strain S288C (ATCC accession number 26108) is modified as described
25 above for use in the method of the invention. The DNA sequence of one unsecreted invertase gene (SUC2) is set forth in SEQ ID NO:1. The original cloning of the SUC2 gene is set forth in M. Carlson et al., Cell 28, 145-154 (1982) and M. Carlson et al., Mol. Cell. Biol. 3, 439-447 (1983), incorporated herein by
30 reference. The DNA sequence of the SUC2 invertase gene is available from

GenBank, having the sequence name YSCSUC2.GB_PL and accession numbers VO1311 and K00540. The SUC2 gene may also be isolated from yeast DNA using the GenBank sequence as the basis for constructing oligonucleotides for use in the polymerase chain reaction.

5 In accordance with the method of the invention, the nonsecreted yeast invertase gene is inserted into a suitable yeast expression vector. Numerous yeast expression vectors are known, for example, the YEp24 expression plasmid having ATCC accession number 37051 and GenBank sequence name YEP24.VEC; and the YRp17 expression plasmid having ATCC accession number 37078 and GenBank
10 sequence name YRP17.VEC. An appropriate yeast expression vector for use in the present invention will contain a suitable yeast promoter and transcription terminator, for example the ADH1 promoter and transcription terminator as described in G. Ammerer, Methods in Enzymology 101, 192-201 (1983), incorporated herein by reference. Preferably, the promoter and transcription terminator are derived from
15 the plasmid AAH5 described in Ammerer. The yeast expression vector will also contain a yeast origin of replication, preferably one which allows extrachromosomal replication; a selectable marker gene for selection of yeast transformants; an *Escherichia coli* origin of replication; and one or more *E. coli* drug resistance genes for selection of *E. coli* transformants. Any yeast origin of replication may be used,
20 so long as it is capable of initiating DNA replication in yeast. Several yeast origins of replication are known, for example, the 2 μ origin, the autonomous replicating sequences (ARS) plus centromeres (CEN elements), and the like. Preferably, the 2 μ yeast origin of replication is used. Similarly, any yeast selectable marker gene may be used, so long as it allows growth only of the desired yeast transformants. Many
25 such yeast selectable marker genes are known; for example URA3, TRP1, and LEU2. Preferably, the TRP1 yeast selectable marker gene is used in the method of the invention. Any *E. coli* origin of replication may be used, so long as it is capable of initiating DNA replication in *E. coli*. Many *E. coli* origins of replication are known, for example pMB1, colE1, and F. Preferably, the pUC *E. coli* origin
30 of replication is used. Any *E. coli* drug resistance gene may be used, so long as it

is capable of allowing growth only of the desired *E.coli* transformants. Many such *E.coli* drug resistance genes are known, for example the ampicillin resistance gene, the chloramphenicol resistance gene, and the tetracycline resistance gene. Preferably, the ampicillin resistance gene is used in the method of the invention.

5 In accordance with the method of the invention, the yeast expression vector containing the nonsecreted yeast invertase gene is ligated to a mammalian cDNA library using known methods. Numerous methods for generating mammalian cDNA libraries are known. For example, poly A⁺ mRNA may be isolated from mammalian cells, purified and the corresponding cDNAs may be synthesized using
10 reverse transcriptase, which is commercially available. Any kind of mammalian cell may be used as a source of mRNA. For example, peripheral blood cells may be used, or primary cells may be obtained from an organ, a developing embryo, a mature animal, a tissue, and the like, and such cells may be the source of mRNA for generation of a cDNA library. Cell lines such as the well known Chinese
15 Hamster Ovary (CHO), COS monkey cell line, the Balb/c 3T3 murine cell line, lymphoid cell lines such as SP 2/0, hybridoma cell lines, and the like, may also be used as a source of mRNA for production of cDNA to construct a library in accordance with the method of the invention.

 In the next step of the method of the invention, the ligated mammalian
20 cDNA-yeast invertase gene is transformed into *E.coli* and recombinants isolated by selecting for drug resistance corresponding to the *E.coli* drug resistance gene on the yeast expression plasmid. Plasmid DNA is then isolated from about one million or more drug resistant *E.coli* recombinants and transformed into yeast strain which does not contain an invertase gene. Many such yeast strains are known, for
25 example the YT455 strain (C. Kaiser et al., Mol. Cell.Biol. 6, 2382-2391 (1986), and yeast strain DA2100, having ATCC accession number 62625. Alternatively, a yeast strain which contains an invertase gene may be manipulated to delete that gene, for example, using methods set forth in Current Protocols in Molecular Biology, Eds. F.M. Ausubel, et al., Jolin Wiley & Sons (1990) in *Saccharomyces cerevisiae* chapters 13.2 and 13.3; or in F. Sherman et al., Methods in Yeast
30

Genetics, Cold Spring Harbor Laboratory (Cold Spring Harbor, N.Y. 1979), both of which are incorporated herein by reference.

In accordance with the method of the invention, yeast recombinants are then selected using a selection pressure corresponding to the yeast selectable marker
5 gene. The yeast recombinants are then collected in pools of about 100,000 to one million transformants and plated on yeast nutrient agar containing only sucrose or only raffinose as the carbon source. This selection will allow growth only of recombinants containing the yeast invertase gene ligated to a mammalian secretory leader sequence. Because invertase deficient yeast may grow on sucrose or
10 raffinose, albeit at a low rate, the invertase selection may be repeated one or more times to maximize the selective pressure applied to the desired recombinants. When the desired number of yeast recombinants are obtained, they are pooled and DNA is isolated and transformed back into *E. coli* for analysis, e.g. by DNA sequencing. Alternatively, DNA may be isolated from individual yeast colonies and analyzed.

15 Novel mammalian secretory leader sequences obtained as described above are purified and used to screen a second cDNA library in the next step of the method of the invention. The second cDNA library is constructed in such a way as to contain full-length cDNAs, using known methods such as those described in Current Protocols in Molecular Biology, chapters 5.5 and 5.6 and in Molecular
20 Cloning, A Laboratory Manual, Second Edition, J. Sambrook, et al., Cold Spring Harbor Laboratory Press (New York, 1989), chapter 8. The full-length cDNAs in the second cDNA library are then ligated to a mammalian expression vector such as the pED vector (Kaufman et al., *Nucleic Acids Res.* 19, 4484-4490 (1991); pEEF-BOS (Mizushima et al., *Nucleic Acids Res.* 18, 5322 (1990); pXM, pJL3 and
25 pJL4 (Gough et al., *EMBO J.* 4, 645-653 (1985); and pMT2 (derived from pMT2-VWF, ATCC accession number 67122, see PCT/US87/00033). The second cDNA library which has been ligated to the mammalian expression vector is transformed into *E. coli*. The library may be screened by hybridization using known screening methods. Alternatively, plasmid DNA is isolated from the transformants for
30 screening by hybridization or using PCR. When screened using PCR, the following

general screening protocol may be followed: the cDNA clone containing the novel leader sequence is sequenced, and appropriate oligonucleotide primers are designed. From about one million *E. coli* transformants, pools of about 100,000 transformants are obtained by spreading pools of 10,000 transformants onto 150 mm plates and
5 replicating the pool onto filters. Plasmid DNA is isolated from each pool and PCR is performed using the oligonucleotide primers based on the novel leader sequence. Specific DNA sequences are detected, for example, by gel electrophoresis of the DNA with or without hybridization. Each of the pools is similarly analyzed, and positive pools are subdivided and purified by hybridizing radioactive
10 oligonucleotides directly to the filters as described in Chapter 6 of Current Protocols in Molecular Biology and in Chapter 1 of Molecular Cloning: A Laboratory Manual.

Using the method set forth above, novel full-length mammalian cDNA clones may be isolated and expressed in transient expression systems such as COS
15 cells grown in a culture medium suitable for growth of cells and production of protein. The novel full-length cDNA clones may also be expressed in stable expression systems such as Chinese hamster ovary cells grown in a culture medium suitable for growth of cells and production of protein. In this way the novel secreted and extracellular proteins of the invention encoded by the mammalian
20 cDNAs are produced which may then be assayed for biological activity in a variety of *in vitro* assays. In addition to detecting novel secreted proteins the method of the invention also detects and allows isolation of integral membrane proteins for example, receptors, and of proteins which transverse the endoplasmic reticulum to localize in intracellular organelles. The novel secreted proteins produced in
25 accordance with the invention may be purified using known methods. For example, the novel secreted protein is concentrated using a commercially available protein concentration filter, such as an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can
30 be employed, for example, a matrix or substrate having pendant diethylaminoethyl

(DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred (e.g., S-Sepharose® columns). The purification of the novel secreted protein from culture supernatant may also include one or more column steps over such affinity resins as lectin-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; or by hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or by immunoaffinity chromatography. Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the novel secreted protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The novel secreted protein thus purified is substantially free of other mammalian proteins.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Jacobs, Kenneth
- (ii) TITLE OF INVENTION: A NOVEL METHOD FOR ISOLATING CYTOKINES
AND OTHER SECRETED PROTEINS
- (iii) NUMBER OF SEQUENCES: 1
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Legal Affairs, Genetics Institute, Inc.
 - (B) STREET: 87 CambridgePark Drive
 - (C) CITY: Cambridge
 - (D) STATE: MA
 - (E) COUNTRY: USA
 - (F) ZIP: 02140
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: McDaniels, Patricia A.
 - (B) REGISTRATION NUMBER: 33,194
 - (C) REFERENCE/DOCKET NUMBER: GI 5200
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 617-876-1170
 - (B) TELEFAX: 617-876-5851

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1542 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Saccharomyces cerevisiae
- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: SUC2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCAATGACAA ACGAAACTAG CGATAGACCT TTGGTCCACT TCACACCCAA CAAGGGCTGG 60
ATGAATGACC CAAATGGGTT GTGGTACGAT GAAAAAGATG CCAAATGGCA TCTGTACTTT 120

CAATACAACC CAAATGACAC CGTATGGGGT ACGCCATTGT TTTGGGGCCA TGCTACTTCC	180
GATGATTTGA CTAATTGGGA AGATCAACCC ATTGCTATCG CTCCCAAGCG TAACGATTCA	240
GGTGCTTTCT CTGGCTCCAT GGTGGTTGAT TACAACAACA CGAGTGGGTT TTTCAATGAT	300
ACTATTGATC CAAGACAAAG ATGCGTTGCG ATTTGGACTT ATAACACTCC TGAAAGTGAA	360
GAGCAATACA TTAGCTATTC TCTTGATGGT GGTACACTT TTAAGAATA CCAAAGAAGC	420
CCTGTTTTAG CTGCCAACTC CACTCAATTC AGAGATCCAA AGGTGTTCTG GTATGAACCT	480
TCTCAAAAAT GGATTATGAC GGCTGCCAAA TCACAAGACT ACAAATTGA AATTTACTCC	540
TCTGATGACT TGAAGTCCTG GAAGCTAGAA TCTGCATTG CCAACGAAGG TTTCTTAGGC	600
TACCAATACG AATGTCCAGG TTTGATTGAA GTCCCAACTG AGCAAGATCC TTCCAAATCT	660
TATTGGGTCA TGTATTATTC TATCAACCCA GGTGCACCTG CTGGCGGTTT CTTCAACCAA	720
TATTTTGTG GATCCTTCAA TGGTACTCAT TTTGAAGCGT TTGACAATCA ATCTAGAGTG	780
GTAGATTTTG GTAAGGACTA CTATGCCTTG CAACTTTCT TCAACACTGA CCCAACCTAC	840
GGTTCAGCAT TAGGTATTGC CTGGGCTTCA AACTGGGAGT ACAGTGCCTT TGTCCCAACT	900
AACCCATGGA GATCATCCAT GTCTTTGGTC CGCAAGTTT CTTTGAACAC TGAATATCAA	960
GCTAATCCAG AGACTGAATT GATCAATTTG AAAGCCGAAC CAATATTGAA CATTAGTAAT	1020
GCTGGTCCCT GGTCTCGTTT TGCTACTAAC ACAACTCTAA CTAAGGCCAA TTCTTACAAT	1080
GTCGATTTGA GCAACTCGAC TGGTACCCTA GAGTTTGAGT TGGTTTACGC TGTTAAACACC	1140
ACACAAACCA TATCCAAATC CGTCTTTGCC GACTTATCAC TTTGGTTCAA GGGTTTAGAA	1200
GATCCTGAAG AATATTTGAG AATGGGTTTT GAAGTCAGTG CTCTTCCTT CTTTTTGGAC	1260
CGTGGTAACT CTAAGGTCAA GTTTGTCAAG GAGAAGCCAT ATTCACAAA CAGAATGTCT	1320
GTCAACAACC AACCATTCAA GTCTGAGAAC GACCTAAGTT ACTATAAAGT GTACGGCCTA	1380
CTGGATCAAA ACATCTTGGA ATTGTACTTC AACGATGGAG ATGTGGTTTC TACAAATACC	1440
TACTTCATGA CCACGGTAA CGCTCTAGGA TCTGTGAACA TGACCACTGG TGTCGATAAT	1500
TTGTTCTACA TTGACAAGTT CCAAGTAAGG GAAGTAAAT AG	1542

CLAIMS

I claim:

1. A method for isolating a cDNA encoding a novel secreted mammalian protein which comprises:
 - a) constructing a cDNA library from mammalian cellular RNA;
 - b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
 - c) transforming the ligated DNA into *E. coli*;
 - d) isolating plasmid DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed *E. coli* of step c);
 - e) transforming the DNA of step d) into a yeast cell which does not contain an invertase gene;
 - f) selecting yeast cells capable of growth on sucrose or raffinose;
 - g) purifying DNA from step f) which contains a novel mammalian leader sequence ligated to yeast invertase;
 - h) screening a second cDNA library to detect full-length cDNAs which contain the novel mammalian leader sequence of step g); and
 - i) isolating the full-length cDNA of step h).
2. A cDNA encoding a secreted protein isolated using the method of:
 - a) constructing a cDNA library from mammalian cellular RNA;
 - b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
 - c) transforming the ligated DNA into *E. coli*;
 - d) isolating plasmid DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed *E. coli* of step c);
 - e) transforming the DNA of step d) into a yeast cell which does not contain an invertase gene;

- f) selecting yeast cells capable of growth on sucrose or raffinose;
 - g) purifying DNA from step f) which contains a novel mammalian leader sequence ligated to yeast invertase;
 - h) screening a second cDNA library to detect full-length cDNAs which contain the novel mammalian leader sequence of step g); and
 - i) isolating the full-length cDNA of step h).
3. A secreted protein isolated by the steps of:
- a) constructing a cDNA library from mammalian cellular RNA;
 - b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
 - c) transforming the ligated DNA into *E. coli*;
 - d) isolating plasmid DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase gene from the transformed *E. coli* of step c);
 - e) transforming the DNA of step d) into a yeast cell which does not contain the invertase gene;
 - f) selecting yeast cells capable of growth on sucrose or raffinose;
 - g) purifying DNA from step f) which contains a novel mammalian leader sequence ligated to yeast invertase;
 - h) screening a second cDNA library to detect full-length cDNAs which contain the novel mammalian leader sequence of step g);
 - i) isolating the full-length cDNA of step h);
 - j) expressing the full-length cDNA of step i) in a host cell grown in a suitable culture medium; and
 - k) isolating the protein expressed in step j) from the culture medium.

4. A method for screening for a cDNA encoding a novel secreted mammalian protein which comprises:

- a) constructing a cDNA library from mammalian cellular RNA;
- b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
- c) transforming the DNA of step b) into a yeast cell which does not contain an invertase gene;
- d) selecting transformed yeast cells from step c) which are capable of growth on sucrose or raffinose;
- e) purifying DNA from the yeast cells of step d);
- f) analyzing the DNA obtained from step e) to determine its sequence and to determine whether it contains a novel sequence;
- g) preparing a second cDNA library from mammalian cellular RNA and screening said second cDNA library to detect full-length cDNAs which contain the novel sequence of step f); and
- h) isolating the full-length cDNA of step g) wherein the isolated cDNA encodes a putative secreted mammalian protein.

5. The method of claim 4, comprising the following additional steps:

- 1) transforming the ligated DNA into bacteria;
- 2) isolating DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed bacteria of step 1);

wherein said additional steps are performed after step b) and before step c), and wherein the DNA isolated in step 2) is used for the transformation in step c).

6. The method of claim 5 wherein said bacteria is *E. coli*.

7. A cDNA encoding a novel secreted protein, wherein said cDNA is isolated by a method comprising:

- a) constructing a cDNA library from mammalian cellular RNA;
- b) ligating said cDNA library to a DNA encoding a nonsecreted

- yeast invertase;
- c) transforming the DNA of step b) into a yeast cell which does not contain an invertase gene;
 - d) selecting transformed yeast cells from step c) which are capable of growth on sucrose or raffinose;
 - e) purifying DNA from the yeast cells of step d);
 - f) analyzing the DNA obtained from step e) to determine its sequence and to determine whether it contains a novel sequence;
 - g) preparing a second cDNA library from mammalian cellular RNA and screening said second cDNA library to detect full-length cDNAs which contain the novel sequence of step f); and
 - h) isolating the full-length cDNA of step g) wherein the isolated cDNA encodes a putative secreted mammalian protein.

8. The cDNA of claim 7, wherein the method comprises the following additional steps:

- 1) transforming the ligated DNA into bacteria;
- 2) isolating DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed bacteria of step 1);

wherein said additional steps are performed after step b) and before step c), and wherein the DNA isolated in step 2) is used for the transformation in step c).

9. The cDNA of claim 8 wherein in said method said bacteria is *E. coli*.

10. A secreted mammalian protein isolated by a method comprising:

- a) constructing a cDNA library from mammalian cellular RNA;
- b) ligating said cDNA library to a DNA encoding a nonsecreted yeast invertase;
- c) transforming the DNA of step b) into a yeast cell which does not

contain an invertase gene;

- d) selecting transformed yeast cells from step c) which are capable of growth on sucrose or raffinose;
- e) purifying DNA from the yeast cells of step d);
- f) analyzing the DNA obtained from step e) to determine its sequence and to determine whether it contains a novel sequence;
- g) preparing a second cDNA library from mammalian cellular RNA and screening said second cDNA library to detect full-length cDNAs which contain the novel sequence of step f);
- h) isolating the full-length cDNA of step g) wherein the isolated cDNA encodes a putative secreted mammalian protein;
- i) expressing the full-length cDNA of step h) in a host cell grown in a suitable culture medium; and
- j) isolating the protein expressed in step i) from the culture

medium.

11. The protein of claim 10, wherein the method comprises the following additional steps:

- 1) transforming the ligated DNA into bacteria;
- 2) isolating DNA containing mammalian cDNA ligated to the DNA encoding the nonsecreted yeast invertase from the transformed bacteria of step 1);

wherein said additional steps are performed after step b) and before step c), and wherein the DNA isolated in step 2) is used for the transformation in step c).

12. The protein of claim 11 wherein in said method said bacteria is *E. coli*.

INTERNATIONAL SEARCH REPORT

International Application No
PC1/US 96/05727

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/10 C12N9/26 C12N15/62

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 607 054 (HONJO TASUKU ;ONO PHARMACEUTICAL CO (JP)) 20 July 1994 see the whole document ---	1-12
Y	US,A,4 914 025 (MANOIL COLIN ET AL) 3 April 1990 see column 6, line 1 - column 7, line 21; claims 1,4 --- -/--	1-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

29 November 1996

Date of mailing of the international search report

13.12.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Chambonnet, F

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/05727

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SCIENCE, vol. 315, 16 January 1987, LANCASTER, PA US, pages 312-317, XP000610609 KAISER, C. A. ET AL.: "Many random sequences functionally replace the secretion signal sequence of yeast invertase" see page 312, column 2, paragraph 3 - page 313, column 1, paragraph 1; figure 1 ---	1-12
A	JOURNAL OF CELLULAR BIOCHEMISTRY, vol. suppl, no. 21a, 10 March 1995, MD US, page 19 XP002019850 JACOBS, K. ET AL.: "A novel secreted method for isolating eukaryotic cDNA clones encoding secreted protein" see abstract C1-207 ---	1
O,A	& Keystone Symposium on dendritic cells: Antigen presenting cells of T and B lymphocytes. Taos, USA, March 10-16 1995 ---	
A	EP,A,0 244 042 (GIST BROCADES NV) 4 November 1987 see the whole document ---	1
E	US,A,5 536 637 (JACOBS KENNETH) 16 July 1996 see the whole document -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/05727

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0607054	20-07-94	CA-A- 2113363 JP-A- 6315380 US-A- 5525486	15-07-94 15-11-94 11-06-96
US-A-4914025	03-04-90	NONE	
EP-A-0244042	04-11-87	AT-T- 110111 DE-D- 3750378 DE-T- 3750378 ES-T- 2061482 IE-B- 64916 JP-A- 63039586 US-A- 5037760 US-A- 5212070	15-09-94 22-09-94 19-01-95 16-12-94 20-09-95 20-02-88 06-08-91 18-05-93
US-A-5536637	16-07-96	NONE	